



24 March 2021

The Royal Australian and New Zealand College of Radiologists Statement on Vaccine Induced Adenopathy

Recent literature from the United States has indicated that some individuals own immune response post COVID vaccination (Pfizer and Moderna) may manifest as unilateral axillary lymphadenopathy related to the vaccinated arm.^{1, 2}

Such responses have not been reported with the Astra-Zeneca vaccine thus far.

Unilateral axillary lymphadenopathy may present as a palpable lump in the axilla clinically. It may inadvertently be seen on routine breast imaging. Clinical history (correlating timing and site of vaccination) and examination should guide appropriate management, including follow up and further investigation, if required. Most vaccination related lymphadenopathy will resolve within 1-2 weeks, some lasting up to 6 weeks. If lymphadenopathy persists, further investigations should be undertaken to rule out alternative conditions.

The Royal Australian and New Zealand College of Radiologists does not recommend delays to vaccination nor any breast imaging. Women are advised to have their screening mammograms before being vaccinated or six weeks after. Patients with unilateral axillary lymphadenopathy should be managed on a case-by-case basis.

References

¹ [Unilateral axillary Adenopathy in the setting of COVID-19 vaccine - ScienceDirect](#)

² [Mitigating the Impact of Coronavirus Disease \(COVID-19\) Vaccinations on Patients Undergoing Breast Imaging Examinations: A Pragmatic Approach : American Journal of Roentgenology : Ahead of Print \(AJR\) \(ajronline.org\)](#)

Please direct any queries to the Faculty of Clinical Radiology at fcr@ranzcr.edu.au

Approved by:
RANZCR President and Dean of the Faculty of Radiology

Date of approval:
24 March 2021

